



Goat

Purina® Goat Nutrition Program


Purina® Goat Grower 16 Plus Up DQ.0015



Complete feed for the growth and development of goats

Purina® Goat Grower 16 Plus Up DQ.0015 is a pelleted complete feed formulated for the optimum growth and development of goats. Whether you raise meat goats or dairy goats, Purina® Goat Grower 16 Plus Up DQ.0015 feed is designed to meet their exacting needs. Purina® products deliver the nutrition and performance you expect. Try Purina® Goat Grower 16 Plus Up DQ.0015 feed and see the difference it can make in your goats' overall growth and health.

FEATURE	BENEFIT
Nutritionally Complete	Helps provide the proper balance of high-quality proteins, vitamins, minerals and other nutrients Built in roughage source helps minimize clean-up from wasted or uneaten forage
Palatable, Premium Ingredients	Consistent quality helps assure high palatability and supports performance
Pelleted	Helps reduce sorting, easy to handle
Organic Trace Minerals	Provides critical highly-available minerals that help support immune function and allows goats to reach their full potential
Urinary Acidifiers including Ammonium Chloride and/or Ammonium Sulfate	Helps reduce the likelihood of urinary calculi
Yeast Culture	Helps maintain rumen health and digestive efficiency
Options	Non-medicated option available (depending on plant)



FEED GREATNESS®



Goat

Purina® Goat Nutrition Program

Purina® Goat Grower 16 Plus Up DQ.0015



Guaranteed Analysis:

ACTIVE DRUG INGREDIENTS

Decoquinatate0.0015 %
(6.8 mg/lb)

NUTRIENT	LEVEL
Crude Protein, (Min)	16.00%
This includes not more than 1.5% equivalent crude protein from non-protein nitrogen	
Crude Fat, (min).....	2.00%
Crude Fiber, (max)	16.00%
Acid Detergent Fiber (ADF) (max).....	20.00%
Calcium (Ca), (min)	1.00%
Calcium (Ca), (max)	1.50%
Phosphorus (P), (min)	0.50%
Salt (NaCl), (min).....	0.50%
Salt (NaCl), (max).....	1.00%
Sodium (Na), (max)	0.65%
Copper (Cu), (min)	20.00 ppm
Copper (Cu), (max)	27.60 ppm
Selenium (Se), (min).....	0.30 ppm
Vitamin A, (min).....	5,000 IU/lb

Ingredients*:

Wheat Middlings, Wheat Red Dog, Ground Soybean Hulls, Corn Distillers Dried Grains with Solubles, Cane Molasses, Calcium Carbonate, Dehydrated Alfalfa Meal, Dehulled Soybean Meal, Ground Corn, Ammonium Chloride, Salt, Yeast Culture, Colored with Iron Oxide, Vitamin A Supplement, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Vitamin D₃ Supplement, Cobalt Glucoheptonate, Vitamin E Supplement, Zinc Oxide, Cobalt Carbonate, Manganese Sulfate, Ethylenediamine Dihydroiodide, Zinc Sulfate, Sodium Selenite, Basic Copper Chloride.

*Ingredients differ by manufacturing plant.

Net WT. 50 lbs (22.67 kg)
Item Number 3009591-201/206
Formula Code: 55LX

Directions:

Feed at a rate of 1.67 lbs to goats/kid weighing 50 lbs to provide 22.7 mg decoquinatate per 100 lb of bodyweight (0.5 mg/kg) per day. Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard.

Warning: Do not feed to goats producing milk for food.

Important:

1. Start kids on colostrum milk from birth to 3 days of age; feed Land O'Lakes® Doe's Match® Kid Milk Replacer or Purina® Goat Kid Milk Replacer from 2 days until 4-8 weeks of age.
2. If kids have not been creep fed or they were purchased, then they should be placed on grass hay on a free-choice basis after arrival.
3. When making a ration change, allow 7 to 10 days for animals to adjust to the new ration.
4. Keep a constant supply of complete feed available. Do not let fine material accumulate in feeders.
5. Provide adequate bunk space for each animal. Bunks should be well-protected and well-managed to prevent feed from becoming wet or moldy.
6. Provide a source of fresh, clean water at all times.
7. Feed Purina® Goat Mineral free-choice.
8. Consult your veterinarian for the recommended health program for your local area. This includes internal and external parasite control. In groups of goats there are certain animals that experience chronic bloat, or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption and poorly managed feeding practices, can increase the incidence of bloat in all goats. If bloating occurs, the above management practices should be reviewed and your veterinarian consulted.

Caution:

Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death.

This product contains copper and should not be fed to sheep. Bentonite should not be used in decoquinatate feeds.

USE ONLY AS DIRECTED.



FEED GREATNESS®