



Show Cattle



Purina® Honor® Show Feed

Fitters Edge® 13 DX

Honor® Show Feed Fitter's Edge® feed is a coarse-textured sweet feed. The versatility of this ration makes this the “tried-and-true” staple of the Honor® Show lineup. Providing flexibility from start to finish, Fitter's Edge® can be fed as a complete ration to grow and develop show cattle. Formulated with 13% protein and moderate levels of energy and fiber. Can also be used to help “stretch one out” without losing rib shape and condition. Get your best animals on Honor® Show Feed Fitter's Edge® feed today so they can reach their full potential.

FEATURE

BENEFIT

Zinpro® Availa® 4 Minerals

- Balanced combination of zinc, manganese, copper and cobalt.
- Provides critical, highly bioavailable minerals that help support immune function and hoof integrity.

Peloton® Yeast feed additive

- Thermal stable yeast encapsulation maintains effectiveness of yeast during pelleting.
- Optimized feed digestibility, supporting feed intake, and animal performance from *Saccharomyces cerevisiae* yeast.
- PMI Fulfill® Feed Additive Mannans (MOS) and beta-glucans support gut integrity and immune-competence.

Research Proven

Feed it with total confidence, backed by Purina's legendary tradition of fundamental and applied research. Along with thousands of winning entries at local and national shows.

Medicated with Deccox® (decoquinate)

For the prevention of coccidiosis. No withdrawal period prior to slaughter.



FEED GREATNESS®



Fitters Edge® 13 DX

Guaranteed Analysis:

ACTIVE DRUG INGREDIENTS

Decoquinatate0.00250%
(11.35 mg/lb)

For the prevention of coccidiosis in cattle (ruminating and non-ruminating calves and veal calves) by Eimeria bovis and E. zuernii

NUTRIENT	LEVEL
Crude protein, min	13.00%
Crude fat, min	2.50%
Crude fiber, max	15.00%
Calcium (Ca), min	0.58%
Calcium (Ca), max	1.08%
Phosphorous (P), min	0.40%
Salt (NaCl), min	0.15%
Salt (NaCl), max	0.65%
Sodium (Na), max	0.51%
Potassium (K), min	0.65%
Selenium (Se), min	0.44 ppm
Selenium (Se), max	0.53 ppm
Vitamin A, min	3,520 IU/lb

Directions:

Feed at the rate of 2 lbs per 100 lbs of bodyweight to provide 22.7 mg Decoquinatate per 100 pounds of body weight (0.5 mg/kg) per day. Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard.

Follow these management practices:

1. Monitor body condition of cattle on a regular basis.
2. Recommended feeding rates should be adjusted depending on cattle body condition.
3. When making a ration change, allow 7 to 10 days for animals to adjust to the new ration.
4. Provide adequate feeder space for each animal.
5. Provide mineral and salt free choice.
6. Provide fresh, clean water near the feeding area.

In groups of cattle, there are certain animals that experience chronic bloat or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption and poorly managed feeding practices can increase the incidence of bloat in all cattle. If bloat occurs, please review feed management practices.


Caution: Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals.



High Octane® supplements are specifically designed for livestock show animals. These supplements are part of the Honor® Show lineup of livestock show products.



OUR WINNING PHILOSOPHY



Check your current price list for availability in your region, or contact your local Purina Retailer or Representative for more information.

For more information on this product, visit our website at www.purinamills.com or call Purina Customer Service at 1-800-227-8941.

Visit our website at www.purinamills.com

Product Code: 3011976-506 / 39ZS - Flaked Corn
3011975-506 / 39ZY - Cracked Corn
*product version depends on regional availability



FEED GREATNESS®